

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1,2-Bis[2-[(4-methoxybenzylidene)-amino]phenyl]disulfane

Wei Xin

Kingfa Scientific & Technological Coporation Limited, Guang Zhou, China 510663, People's Republic of China

Correspondence e-mail: xinw0408@163.com


Received 15 April 2013; accepted 28 April 2013

Key indicators: single-crystal X-ray study; $T = 298$ K; mean $\sigma(\text{C}-\text{C}) = 0.004$ Å; R factor = 0.051; wR factor = 0.139; data-to-parameter ratio = 14.0.

The asymmetric unit of the title compound, $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_2\text{S}_2$, contains one-half molecule, which is completed by twofold rotation symmetry with the twofold axis passing through the mid-point of the central S—S bond. The planes of the two benzene rings linked by the disulfide chain form a dihedral angle of $76.1(1)^\circ$, while the planes of the two benzene rings in the benzylideneaniline fragment form a dihedral angle of $48.9(1)^\circ$. The crystal packing exhibits no significantly short intermolecular contacts.

Related literature

For the crystal structures of related compounds, see: İde *et al.* (1997); Ozbey *et al.* (1998); He *et al.* (2011); Wang *et al.* (2011).



Experimental

Crystal data

 $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_2\text{S}_2$ $M_r = 484.61$ Orthorhombic, $Pbcn$ $a = 10.2657(11)$ Å $b = 13.0675(13)$ Å $c = 18.3415(15)$ Å $V = 2460.5(4)$ Å³ $Z = 4$ Mo $K\alpha$ radiation $\mu = 0.25$ mm⁻¹ $T = 298$ K $0.26 \times 0.22 \times 0.17$ mm

Data collection

Bruker SMART APEX CCD area-detector diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

 $T_{\min} = 0.939$, $T_{\max} = 0.960$

9487 measured reflections

2163 independent reflections

1363 reflections with $I > 2\sigma(I)$ $R_{\text{int}} = 0.094$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.051$ $wR(F^2) = 0.139$ $S = 0.95$

2163 reflections

155 parameters

H-atom parameters constrained

 $\Delta\rho_{\text{max}} = 0.37$ e Å⁻³ $\Delta\rho_{\text{min}} = -0.28$ e Å⁻³

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL*.

The author acknowledges the financial support of the Kingfa Scientific & Technological corporation Ltd.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5405).

References

- Bruker (2007). *SMART* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- He, Q.-P., Dai, L. & Tan, B. (2011). *Acta Cryst.* **E67**, o3240.
- İde, S., Öztaş, G., Ancin, N. & Tüzün, M. (1997). *Acta Cryst.* **C53**, 376–378.
- Ozbey, S., Temel, A., Ancin, N., Öztas, S. G. & Tuzun, M. (1998). *Z. Kristallogr. New Cryst. Struct.* **213**, 207–208.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Wang, Y., Shi, S., Han, Y. & Wei, G.-D. (2011). *Acta Cryst.* **E67**, o3364.

supplementary materials

Acta Cryst. (2013). E69, o896 [doi:10.1107/S1600536813011598]

1,2-Bis{2-[(4-methoxybenzylidene)amino]phenyl}disulfane

Wei Xin

Comment

As a contribution to structural study of diaminodiphenyl disulfides (Ide *et al.*, 1997; Wang *et al.*, 2011; He *et al.*, 2011), we present here the crystal structure of the title compound, (I).

In (I) (Fig. 1), the bond lengths and angles are normal and correspond to those observed in related compounds (Ide *et al.*, 1997; Ozbey *et al.*, 1998; He *et al.*, 2011). The molecule has crystallographic twofold rotation symmetry with the twofold axis passing through the midpoint of the central S—S bond. Two benzene rings connected through disulfide chain form a dihedral angle of 76.1 (1)°. Two benzene rings in two benzylideneaniline fragments form the dihedral angles of 48.9 (1)°. The crystal packing exhibits no significantly short intermolecular contacts.

Experimental

4-Methoxybenzaldehyde (2 mmol) in ethanol (10 ml) was added to a solution of 2,2'-diaminodiphenyl disulfide (1 mmol) in ethanol (20 ml). The solution was heated to 323 K for 4 h. The reaction mixture was cooled to room temperature and the yellow crystalline product was obtained.

Refinement

All H atoms were placed in geometrically idealized positions (C—H 0.93–0.96 Å) and treated as riding on their parent atoms, with $U_{\text{iso}}(\text{H}) = 1.2\text{--}1.5U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *SMART* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008).




Figure 1

The molecular structure of (I) showing the atomic numbering and 50% probability displacement ellipsoids. Unlabelled atoms are related with the labelled ones by symmetry operation $(-x, y, 1/2 - z)$.

1,2-Bis{2-[(4-methoxybenzylidene)amino]phenyl}disulfane

Crystal data

$C_{28}H_{24}N_2O_2S_2$

$M_r = 484.61$

Orthorhombic, *Pbcn*

$a = 10.2657$ (11) Å

$b = 13.0675$ (13) Å

$c = 18.3415$ (15) Å

$V = 2460.5$ (4) Å³

$Z = 4$

$F(000) = 1016$

$D_x = 1.308$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 1719 reflections

$\theta = 2.5\text{--}22.9^\circ$

$\mu = 0.25$ mm⁻¹

$T = 298$ K

Block, yellow

$0.26 \times 0.22 \times 0.17$ mm

Data collection

Bruker SMART APEX CCD area-detector
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ϕ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

$T_{\min} = 0.939$, $T_{\max} = 0.960$

9487 measured reflections

2163 independent reflections

1363 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.094$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.5^\circ$

$h = -8 \rightarrow 12$

$k = -12 \rightarrow 15$

$l = -17 \rightarrow 21$

Refinement

Refinement on F^2

Least-squares matrix: full

 $R[F^2 > 2\sigma(F^2)] = 0.051$
 $wR(F^2) = 0.139$
 $S = 0.95$

2163 reflections

155 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: inferred from
neighbouring sites

H-atom parameters constrained

 $w = 1/[\sigma^2(F_o^2) + (0.075P)^2]$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$
 $\Delta\rho_{\max} = 0.37 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.28 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.01987 (8)	0.02789 (6)	0.19518 (4)	0.0510 (3)
N1	0.1694 (2)	0.02278 (17)	0.06606 (11)	0.0433 (6)
O1	0.0529 (2)	0.39678 (18)	−0.14776 (12)	0.0757 (7)
C1	0.1483 (3)	−0.0635 (2)	0.18144 (14)	0.0408 (7)
C2	0.1829 (3)	−0.1399 (2)	0.23057 (15)	0.0494 (8)
H2	0.1390	−0.1459	0.2747	0.059*
C3	0.2830 (3)	−0.2067 (2)	0.21360 (16)	0.0574 (8)
H3	0.3062	−0.2575	0.2467	0.069*
C4	0.3491 (3)	−0.1991 (2)	0.14808 (16)	0.0596 (9)
H4	0.4153	−0.2450	0.1369	0.072*
C5	0.3159 (3)	−0.1225 (2)	0.09907 (15)	0.0510 (8)
H5	0.3605	−0.1171	0.0551	0.061*
C6	0.2160 (3)	−0.0532 (2)	0.11509 (14)	0.0417 (7)
C7	0.2471 (3)	0.0735 (2)	0.02534 (14)	0.0468 (7)
H7	0.3359	0.0594	0.0276	0.056*
C8	0.2011 (3)	0.1530 (2)	−0.02483 (13)	0.0451 (7)
C9	0.0684 (3)	0.1658 (2)	−0.03945 (14)	0.0483 (7)
H9	0.0089	0.1192	−0.0205	0.058*
C10	0.0241 (3)	0.2456 (2)	−0.08118 (15)	0.0518 (8)
H10	−0.0646	0.2524	−0.0902	0.062*
C11	0.1108 (3)	0.3167 (2)	−0.11021 (14)	0.0518 (8)
C12	0.2432 (3)	0.3044 (2)	−0.09884 (14)	0.0541 (8)
H12	0.3024	0.3505	−0.1187	0.065*
C13	0.2868 (3)	0.2216 (2)	−0.05696 (14)	0.0539 (8)
H13	0.3758	0.2123	−0.0505	0.065*
C14	0.1355 (4)	0.4714 (3)	−0.1798 (2)	0.0852 (12)

H14A	0.1930	0.4987	−0.1434	0.128*
H14B	0.0836	0.5257	−0.1998	0.128*
H14C	0.1859	0.4404	−0.2180	0.128*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0584 (6)	0.0438 (5)	0.0509 (5)	0.0064 (4)	0.0120 (4)	0.0093 (3)
N1	0.0474 (15)	0.0441 (13)	0.0383 (12)	−0.0016 (11)	0.0033 (11)	0.0047 (11)
O1	0.0769 (17)	0.0664 (16)	0.0839 (16)	−0.0040 (13)	−0.0048 (13)	0.0296 (13)
C1	0.0446 (18)	0.0362 (15)	0.0415 (15)	−0.0036 (13)	0.0016 (13)	−0.0014 (12)
C2	0.064 (2)	0.0450 (18)	0.0395 (15)	−0.0004 (15)	0.0080 (14)	0.0022 (13)
C3	0.069 (2)	0.0495 (19)	0.0536 (18)	0.0127 (17)	0.0051 (16)	0.0100 (15)
C4	0.064 (2)	0.054 (2)	0.061 (2)	0.0174 (16)	0.0045 (16)	0.0008 (16)
C5	0.054 (2)	0.0534 (19)	0.0451 (16)	0.0032 (16)	0.0130 (14)	−0.0001 (14)
C6	0.0428 (17)	0.0428 (16)	0.0395 (15)	−0.0035 (13)	−0.0009 (13)	−0.0001 (12)
C7	0.0447 (18)	0.0508 (17)	0.0449 (15)	−0.0015 (15)	0.0022 (14)	0.0015 (13)
C8	0.053 (2)	0.0492 (17)	0.0330 (14)	−0.0047 (14)	0.0048 (13)	−0.0003 (12)
C9	0.0509 (19)	0.0487 (18)	0.0452 (16)	−0.0089 (15)	0.0002 (14)	0.0031 (13)
C10	0.0499 (19)	0.0558 (19)	0.0496 (17)	−0.0019 (16)	−0.0017 (14)	0.0028 (15)
C11	0.063 (2)	0.0533 (19)	0.0389 (15)	−0.0033 (17)	−0.0038 (15)	0.0040 (14)
C12	0.065 (2)	0.0540 (19)	0.0436 (16)	−0.0157 (17)	0.0081 (15)	0.0101 (14)
C13	0.0511 (19)	0.066 (2)	0.0444 (16)	−0.0081 (16)	0.0066 (14)	0.0069 (15)
C14	0.101 (3)	0.065 (3)	0.090 (3)	−0.011 (2)	0.005 (2)	0.030 (2)

Geometric parameters (\AA , $^\circ$)

S1—C1	1.797 (3)	C7—C8	1.466 (4)
S1—S1 ⁱ	2.0518 (14)	C7—H7	0.9300
N1—C7	1.278 (3)	C8—C13	1.388 (4)
N1—C6	1.423 (3)	C8—C9	1.398 (4)
O1—C11	1.387 (4)	C9—C10	1.372 (4)
O1—C14	1.420 (4)	C9—H9	0.9300
C1—C2	1.390 (4)	C10—C11	1.392 (4)
C1—C6	1.408 (3)	C10—H10	0.9300
C2—C3	1.385 (4)	C11—C12	1.385 (4)
C2—H2	0.9300	C12—C13	1.400 (4)
C3—C4	1.383 (4)	C12—H12	0.9300
C3—H3	0.9300	C13—H13	0.9300
C4—C5	1.388 (4)	C14—H14A	0.9600
C4—H4	0.9300	C14—H14B	0.9600
C5—C6	1.398 (4)	C14—H14C	0.9600
C5—H5	0.9300		
C1—S1—S1 ⁱ	106.46 (9)	C13—C8—C9	117.3 (3)
C7—N1—C6	121.4 (2)	C13—C8—C7	121.3 (3)
C11—O1—C14	117.9 (3)	C9—C8—C7	121.3 (3)
C2—C1—C6	120.1 (3)	C10—C9—C8	121.4 (3)
C2—C1—S1	125.0 (2)	C10—C9—H9	119.3
C6—C1—S1	114.8 (2)	C8—C9—H9	119.3

C3—C2—C1	119.8 (3)	C9—C10—C11	120.6 (3)
C3—C2—H2	120.1	C9—C10—H10	119.7
C1—C2—H2	120.1	C11—C10—H10	119.7
C4—C3—C2	120.9 (3)	C12—C11—O1	125.7 (3)
C4—C3—H3	119.6	C12—C11—C10	119.5 (3)
C2—C3—H3	119.6	O1—C11—C10	114.8 (3)
C3—C4—C5	119.7 (3)	C11—C12—C13	119.1 (3)
C3—C4—H4	120.2	C11—C12—H12	120.5
C5—C4—H4	120.2	C13—C12—H12	120.5
C4—C5—C6	120.7 (3)	C8—C13—C12	122.0 (3)
C4—C5—H5	119.7	C8—C13—H13	119.0
C6—C5—H5	119.7	C12—C13—H13	119.0
C5—C6—C1	118.8 (2)	O1—C14—H14A	109.5
C5—C6—N1	124.4 (2)	O1—C14—H14B	109.5
C1—C6—N1	116.5 (2)	H14A—C14—H14B	109.5
N1—C7—C8	122.2 (3)	O1—C14—H14C	109.5
N1—C7—H7	118.9	H14A—C14—H14C	109.5
C8—C7—H7	118.9	H14B—C14—H14C	109.5

Symmetry code: (i) $-x, y, -z+1/2$.